



ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

CODEN: IJRSFP (USA)

International Journal of Recent Scientific Research  
Vol. 9, Issue, 4(F), pp. 26068-26071, April, 2018

**International Journal of  
Recent Scientific  
Research**

DOI: 10.24327/IJRSR

## Research Article

# THE RELATION BETWEEN HAND GRIP STRENGTH WITH HAND-ANTHROPOMETRIC VARIABLE IN INTER-UNIVERSITY LEVEL SOFTBALL, CRICKET AND HOCKEY PLAYERS

Aswathy V\*

Department of Physical Education, University of Kerala, India

DOI: <http://dx.doi.org/10.24327/ijrsr.2018.0904.1985>

### ARTICLE INFO

#### Article History:

Received 10<sup>th</sup> January, 2018  
Received in revised form 20<sup>th</sup>  
February, 2018  
Accepted 8<sup>th</sup> March, 2018  
Published online 28<sup>th</sup> April, 2018

#### Key Words:

Grip Strength and Anthropometry

### ABSTRACT

The purpose of this study was to compare hand grip strength with hand anthropometric variables of inter-university level Softball, Cricket and hockey players. The study included 45 inter university female softball (15), cricket (15) and hockey (15) players aged 18–25 years. Methods: A total of 11 anthropometric variables and 2 strength variables (Grip strength left, Grip strength right) were selected for the study. The data were collected using standard anthropometric equipment (Gulic tape, sliding caliper, skin fold caliper, and grip dynamometer) which were supplied by reputed Indian and foreign companies and used in the sports science laboratory of Lakshmbai National college of Physical Education. Results: All the three groups showed significant differences on their arm measurements and grip strength variables. The findings of the present study indicated that the grip strength of softball players showed the closest relationship with the arm measurements. Cricket and hockey players correlated second and third respectively. Conclusion: It may be concluded that the grip strength of University level women softball players had significant relationship with most of their arm measurements as compared to cricket and hockey players.

**Copyright © Aswathy V, 2018**, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Softball, Cricket and Hockey are physically demanding sports comprised of several specialism requiring different skills and types of fitness. These sports require excellent eye-to-hand coordination, upper extremity power, hand grip strength and the coordinated movements of the hips, shoulders, arms and wrists. These sports entail bating, throwing, fielding, pitching, running and hitting. Cricket is a game of endurance as well as strength. For bowling, throwing the ball during fielding and batting, use of forearm strength is essential (Koley & Yadav, 2009). Softball is a direct descendant of baseball. It requires speed, strength and endurance (Terbizan, Waldera, Seljevold & Schwigert, 1996). It is a game where the handgrip plays a crucial role. The main basic aspects of the game swing velocity and pitch speed are greatly influenced by the grip strength of the player (Giardina, Leslie, Raridon & Zimmer, 1997). Softball players require a significant amount of upper body muscle balance due to the specificity of underarm activity. Many athletes and coaches believe that the forearm plays a significant role both in hitting and throwing the ball (Brylinskyl & Melanie, 1992). Having good grip strength will allow hockey player to make more solid poke checks, deflect pass and hitting etc.. Hand grip strength is the result of forceful flexion of all finger joints with the maximum voluntary force

that the subject is able to exert under normal bio kinetic conditions (Richards *et al.*, 1996; Bohannon, 1997). Grip strength determines the handedness of an Individual. It is often used as an indicator of the overall physical strength (Foo, 2007). Many studies describe upper extremity characteristics of baseball players (Barnes & Tullos, 1978; King *et al.*, 1969; Torg *et al.*, 1972; Tullos *et al.*, 1972) and reports related to morphological/anthropometric characteristics and hand grip strength of softball players are available (Werner *et al.*, 2006; Koley & Kumaar, 2011). The grip strength is measured in several sports disciplines and its importance to success is clearly identified. The most common method of assessment for grip strength is the use of a handheld dynamometer (Ramkumar & Chittibabu, 2012). The estimation of hand grip strength is of immense importance in sports like wrestling, tennis, badminton, cricket, handball, basketball, hockey, baseball and softball, where sufficient degree of grip strength is necessary to be successful. The word anthropometry is derived from the Greek word `anthros` meaning man, and metre, meaning to measure. Earlier anthropometry was used mainly for racial, Differentiation and personal identification. Anthropometric measurements vary from individual to Individual. The measurement of structure of the body is called Anthropometry. It consists of measuring external environment of the human

\*Corresponding author: **Aswathy V**

Department of Physical Education, University of Kerala, India

body. Due to the high correlation between anthropometric measurements like linear measurement, circumferences, diameters body composition variables etc. and physical performance, the recent trend is to include these variables in the areas related to the strength of the athlete. Anthropometric measurements have revealed correlation between body structure, physical characteristics and sports capabilities. This knowledge of mathematical correlation permits sports physicians to evaluate and to predict performance potentialities on the basis of physical characteristics and specific requirements of the game. Some literature related to the morphological and anthropometric characteristics and handgrip strength of softball players are available (Brylinskyl & Melanie, 1992; Giardina, Leslie, Raridon & Zimmer, 1997; Tsunawake, Tahara, Moji, Muraki, Minowa & Yukawa, 2003; Werner, Deryk, Guido & Brunet, 2006). But, the information related to the correlations of hand-anthropometry and grip strength in softball, cricket and hockey players is scanty. Thus, the purpose of this study was to compare hand grip strength and anthropometric variables of inter-university level Softball, Cricket and hockey players.

## MATERIALS & METHODS

### Subjects

A sample of forty five (N =45) women inter-university level players. The player s’ age ranged from 18 to 25 years. Which includes fifteen each softball, cricket and hockey players, who participated in inter- university competitions of University of Kerala was selected. All the participants were informed about the aim and methodology of the study and they volunteered to participate in this study. Purposive sampling technique was used to select the subjects.

### METHODOLOGY

#### Anthropometric measurements

- **Length measurements:** 1. Acromiale-dactyilion (Arm length) 2. Acromiale-Radiale (upper arm length) 3.Radiale-styilion Radiale (Fore arm length)4. midstyilion-dactyilion (Hand length)
- **Arm girth:** 5. Wrist 6. Fore arm 7. Arm relaxed 8. Arm flexed and tensed
- **Bredth:** 9. Biepicondylarhumerus
- **Skin fold measurements:** 10. Biceps skin fold measurement. 11. Triceps skin fold measurement.

**Table 1** Descriptive statistics of hand grip strength and various anthropometric variables in Soft ball, Cricket and hockey players.

Variables	Group	Left hand				Right hand			
		Mean	SD	Minimum	Maximum	Mean	SD	Minimum	Maximum
AL	Softball	69.15	3.63	60.5	76.5	69.15	3.63	60.5	76.5
	Cricket	69.84	3.52	61.6	74.7	69.8	3.48	61.6	74.7
	Hockey	70.94	2.98	66.5	76.5	71.06	3.08	66.5	76.5
UAL	Softball	28.75	1.26	27	31	28.73	1.27	27	31
	Cricket	29.2	1.52	27	31.3	29.14	1.44	27	31.3
	Hockey	29.32	1.61	26.5	32.5	29.36	1.61	26.5	32.5
FAL	Softball	23.67	0.91	22	25.5	23.74	0.94	22	25.5
	Cricket	23.4	0.96	21.5	25.2	23.42	0.96	21.5	25.2
	Hockey	23.94	1.36	22	26.7	23.88	1.27	22	26.7
HL	Softball	17.41	0.68	16.3	18.6	17.36	0.62	16.3	18.6
	Cricket	17.49	0.98	15.6	19.4	17.49	0.96	15.6	19.4
	Hockey	17.58	0.68	16.7	19.3	17.58	0.65	16.7	19.3
WG	Softball	14.19	1.1	11.6	15.5	14.29	1.12	11.7	16.3
	Cricket	13.93	0.85	12.4	15.5	13.93	0.69	12.2	15
	Hockey	13.88	2.68	5.1	17.5	14.06	2.71	5.2	17.5
FAG	Softball	21.99	1.34	19	24.5	22.26	1.24	19.5	24.5
	Cricket	21.52	1.11	19.1	23.2	21.88	1.21	20	24
	Hockey	22.19	0.86	20.3	23.8	22.57	0.74	21.5	24
ARG	Softball	24.36	2.7	19.5	30.5	25.02	2.64	21	30
	Cricket	23.16	1.66	21	26	24.34	1.8	21	27
	Hockey	24.19	1.75	22	29.1	24.67	2.11	22.3	31.4
AFG	Softball	25.81	2.53	21.5	31	26.25	2.49	22.6	32
	Cricket	25.31	1.95	22.7	28.5	25.94	2.09	22.6	29.5
	Hockey	26.44	1.88	24	31.5	26.97	1.94	24	32
BH	Softball	5.53	0.34	5	6.2	5.59	0.31	5.1	6.2
	Cricket	5.34	0.32	4.7	5.9	5.53	0.39	4.7	6
	Hockey	5.68	0.21	5.2	6	5.8	0.26	5.4	6.3
BSF	Softball	5.88	1.61	2.8	8.6	4.89	1.03	2.6	6.4
	Cricket	5.78	1.79	3.8	10.8	5.28	1.59	2.8	8.2
	Hockey	5.53	1.03	4.2	8	5.02	0.77	3.8	6.2
TSF	Softball	12.5	3.7	7	21.2	13.96	4.03	7.3	20.4
	Cricket	12.18	3.83	6.6	19.4	13.04	4.06	6.8	21.2
	Hockey	11.44	2.78	7	17	12.74	3.27	8	18.4
GS	Softball	20.03	3.19	15.6	26.3	20.72	4.34	15.4	30.3
	Cricket	25.29	3.81	18.6	33.3	26.56	4.45	17.6	35.1
	Hockey	26.66	3.47	19.3	31	28.72	3.94	20.6	34.6

\*Significant at 0.05 level

AL = Arm Length, UAL = Upper arm Length, FAL = Forearm length, HL = Hand Length, WG = Wrist Girth, FAG = Fore arm Girth, ARG = Arm relaxed Girth, AFG = Arm flexed Girth, BH = Biepicondylarhumerus, BSF = Biceps Skin fold, TSF = Triceps Skin fold.

**Strength variables:** 1. Left Grip strength. 2. Right grip strength.

**Statistical analyses**

Standard descriptive statistics (mean ± standard deviation) were determined for directly measured and derived variables. Pearson product moment correlation was applied to establish the correlation of right and left hand grip strength with the variables measured. Data was analysed using SPSS and MS EXCEL 2010. 05 level of probability was used to indicate statistical significance.

**RESULTS**

Table 1 showed that descriptive statistics of hand grip strength and anthropometric variables in inter-university level Softball, Cricket and hockey players.

In the present study, Significant relationships were observed between left grip strength to and wrist girth), arm relaxed girth ( $r = 0.524, p < .05$ ) and arm flexed and tensed- girth ( $r = 0.615, p < .05$ ) of softball players.

Significant relationship was observed between left grip strength and arm length ( $r = .611, p < .05$ ) of cricket players. No significant relationships were observed between left grip strength of hockey players.

Significant relationship were observed between right grip strength and wrist girth ( $r = 0.535, p < .05$ ), Arm relaxed girth ( $r = 0.717, p < .05$ ), arm flexed and tensed girth ( $r = 0.727, p < .05$ ), biepicondylarhumerus ( $r = 0.731, p < .05$ ), biceps Skin fold ( $r = 0.607, p < .05$ ) and triceps Skin fold ( $r = 0.556, < .05$ ) of softball players.

**Table 2** Correlation coefficients of left handgrip strength with hand anthropometric variables ininter-university level Softball, Cricket and hockey players.

Sl no	Variables	Left hand grip strength					
		Softball		Cricket		Hockey	
		R	P	R	P	r	P
1	AL	0.474	0.074	0.611*	0.016	0.124	0.672
2	UAL	0.337	0.220	0.372	0.172	0.262	0.346
3	FAL	0.129	0.647	0.303	0.272	0.001	0.999
4	HL	0.351	0.199	0.553	0.032	0.234	0.401
5	WG	0.680*	0.005	0.459	0.085	0.196	0.483
6	FAG	0.497	0.059	0.575	0.025	0.019	0.947
7	ARG	0.524*	0.045	0.435	0.106	0.240	0.388
8	AFG	0.615*	0.015	0.496	0.060	0.157	0.576
9	BH	0.317	0.250	0.371	0.173	0.030	0.916
10	BSF	0.379	0.163	0.051	0.858	0.367	0.178
11	TSF	0.362	0.186	0.153	0.585	0.157	0.576

AL = Arm Length, UAL = Upper arm Length, FAL = Forearm length, HL = Hand Length, WG = Wrist Girth, FAG = Fore arm Girth, ARG = Arm relaxed Girth, AFG = Arm flexed Girth, BH = Biepicondylarhumerus, BSF = Biceps Skin fold, TSF = Triceps Skin fold.

**Table 3** Correlation coefficients of right handgrip strength with hand anthropometric variables ininter-university level Softball, Cricket and hockey players

Sl no	Variables	Right hand grip strength					
		Softball		Cricket		Hockey	
		R	P	R	P	r	P
1	AL	0.453	0.090	0.724*	0.002	0.256	0.357
2	UAL	0.397	0.143	0.455	0.088	0.387	0.155
3	FAL	0.230	0.410	0.499	0.058	0.186	0.508
4	HL	0.296	0.284	0.560*	0.030	0.410	0.129
5	WG	0.535*	0.040	0.550*	0.033	0.536*	0.039
6	FAG	0.473	0.075	0.763*	0.001	0.055	0.846
7	ARG	0.717*	0.003	0.040	0.887	0.616*	0.015
8	AFG	0.727*	0.002	0.356	0.193	0.470	0.077
9	BH	0.731*	0.002	0.433	0.107	0.206	0.462
10	BSF	0.607*	0.017	0.212	0.449	0.019	0.945

**DISCUSSION**

Soft ball, cricket and hockey are the popular sport and are also an international sport. Anthropometric dimensions and morphological characteristics play an important role in determining the success of an athlete (Keogh, 1999). Quite naturally, the interest in Anthropometric characteristics and body composition of athletes from different competitive sports has increased tremendously over the last decades. All ball games require comprehensive abilities including physical, technical, mental and tactical (Terbizan, Waldera, Seljevod& Schwigert,1996; Werner, Deryk, Guido & Brunt, 2006).

Significant relationship were observed between right grip strength and arm length ( $r = 0.724, p < .05$ ), Hand length ( $r = 0.560, p < .05$ ), Wrist girth ( $r = 0.550, p < .05$ ) and Fore armgirth ( $r = 0.763, p < .05$ ) of cricket players.

Significant relationships were observed between right grip strength to wrist girth ( $r = 0.536, p < .05$ ) and Arm relaxed girth ( $r = 0.616, p < .05$ ) of hockey players.

In the present study all the three groups showed significant differences on their arm measurements and grip strength variables. Thus the hypothesis formulated in the beginning of the study is accepted. Previous studies have been reported that handgrip strength had strong correlation with various

anthropometric characteristics (Benefice & Malina, 1996; Koley & Kaur, 2009; koley & Yadav, 2009; Koley, Kaur & Sandhu, 2009; Kaley, Singh & Sandhu, 2010)

The grip strength of softball players showed the closest relationship with the arm measurements. Cricket and hockey players correlated second and third respectively.

In the present study, small sample size was a limitation which would be taken into account in our future studies.

## CONCLUSIONS

It was concluded that the grip strength of the softball players had significant relationship with most of their arm measurements.

## Acknowledgements

Sincere thanks to coaches who extended their enthusiastic cooperation collecting the necessary data and to all those players without whom this research was not possible. We acknowledge this immense help received from the scholars whose articles are cited and include in reference of this paper. We are also grateful to author/editor/publishers of all those article, journals and books from where the literature of this paper has been reviewed and discussed.

## Reference

- Nath, Surdindhar(1993)“Anthropometry the measurement of body size shape and form”, Friends publications, Delhi
- Jones, Michael Marfell, Olds Tim, Stewart Arthur and Carter Lindsay(2006)“International standard for Anthropometric assessment” Published by the International society for the advancement of Kinanthropometry
- Jurmain, Robert, Nelson Harry, Kilgore Lynn, TrevanthaneWenda(1998)“Essentials of physical Anthropometry” Wordsworth publishing company, 2<sup>nd</sup> Edition
- Sodhi, H S(1991) “Sports Anthropometry:Kinanthropometric” approach Anova publications, Mohali.

## Journals

- McLean, Brian D (1989) “An anthropometric analysis of elite Australian Tracyclists” Journal of Sports Sciences Vol7
- Vishaw, Gaurav (2010) “Anthropometric characteristics, somatotype and body composition of volleyball and basketball players”-*Journal of Physical Education and Sports Mangement.*
- Abraham, George (2010)“anthropometry of Indian athletes”, Indian journal of science and technology, Indian society for education and environment vol. no 12
- Russo, Gualdi *et al.* (1992) “skin fold and body composition of sports participation, *journal sports medicine and physical fitness*” Websites
- Kattel, Bheem P, Fredericks Tycho K, Fernandez Jeffrey E and Lee Dong C. (1996) “The effect of upper-extremity posture on maximum grip strength”
- Blimkie, Cameron J. R, RoachePaul, Hay John TandBar-Or Oded(1988) “Anaerobic power of arms in teenage boys and girls: relationship to lean tissue”<http://link.springer.com/article/10.1007%2FBF01075988#page-1>
- Maglischo, Cheryl W (2013)“Bases of Norms for Cable-Tension Strength Tests for Upper Elementary, Junior High, and Senior High School Girls”, <http://www.tandfonline.com/doi/abs/10.1080/10671188.1968.10616585#.UpIx2sSVMUs>
- Fernandes,(2011) “Correlations between grip strength and anthropometric variables of young adults’ hand” [http://www.scielo.br/scielo.php?pid=S180929502011000200009&script=sci\\_abstract](http://www.scielo.br/scielo.php?pid=S180929502011000200009&script=sci_abstract)
- Clarke, H. Harrison, Petersen Kay H. (2013) “Contrast of Maturational, Structural, and Strength Characteristics of Athletes and Nonathletic10 to15YearsofAge”<http://www.tandfonline.com/doi/abs/10.1080/10671188.1961.10613130#.UpI1IcSVMUt>
- Shyamal, Koley(2011)“A study of arm anthropometric profile in Indian inter-university basketball players”

### How to cite this article:

Aswathy V.2018, The Relation Between Hand Grip Strength with Hand-Nthropometric Variable in Inter-University Level Softball, Cricket and Hockey Players. *Int J Recent Sci Res.* 9(4), pp. 26068-26071.  
DOI: <http://dx.doi.org/10.24327/ijrsr.2018.0904.1985>

\*\*\*\*\*