

ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

CODEN: IJRSFP (USA)

International Journal of Recent Scientific Research  
Vol. 13, Issue, 11(A), pp. 2555-2559, November, 2022

**International Journal of  
Recent Scientific  
Research**

DOI: 10.24327/IJRSR

## Research Article

# RESEARCH ON REASONS FOR NOT ACCEPTING POSTPARTUM FAMILY PLANNING IN FIRST 48 HOURS, IN A TERTIARY CARE HOSPITAL

A.M.R.K. Atapattu\* and P. Gawri Senthilnathan

Department of Obstetrics & Gynaecology, De Soysa Hospital for Women

DOI: <http://dx.doi.org/10.24327/ijrsr.2022.1311.0522>

### ARTICLE INFO

#### Article History:

Received 14<sup>th</sup> October, 2022

Received in revised form 26<sup>th</sup> October, 2022

Accepted 15<sup>th</sup> November, 2022

Published online 28<sup>th</sup> November, 2022

#### Keywords:

Post Partum Family Planning, Subdermal Implant, Post-Partum LRT, Maternal & Perinatal Death, Interval Contraception.

### ABSTRACT

**Introduction:** Postpartum family planning (PPFP) is an essential service to prevent unwanted pregnancies, to keep birth spacing, thus preventing poor maternal, perinatal, and neonatal health outcomes in subsequent pregnancies. **Objectives:** To identify the reasons for not accepting PPFP and actual percentage of PPFP in the first 48 hours as well as the most preferred method of PPFP, most preferred interval contraceptive method in a tertiary care hospital. **Methodology:** The present study is a descriptive cross-sectional study, conducted in all four Obstetric units in De Soysa Hospital for Women, Colombo 08. Inclusion criteria - Postpartum Mothers (within the first 48 hours after delivery) the time from 25<sup>th</sup> of September 2019 until the sample size of 289 is reached. **Results:** 152 (51.5 %) of the participants accepted PPFP. Only 20 (13.3 %) accepted contraceptive methods during the first 48 hours of the postpartum period. Regarding not accepting the PPFP, 53 (37.8%) planned to use in the future (Interval methods), 31 (20.1 %) due to husband's disapproval, 20 (12.1%) due to fear of side effects and 18 (12.9%) wanted to have many children. The most popular PPFP method was Subdermal implant 82 (57.7 %), followed by PPIUD 48 (33.8%) and PP- LRT 12 (8.5%) among participants. **Conclusion:** Approximately 50% of the study population accepted the PPFP and subdermal implant is the commonest mode of contraception. The commonest factor for denying contraception was the expectation of using the Interval contraceptive methods in the future.

Copyright © A.M.R.K. Atapattu 2022, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Family planning (FP) is a very important concept in improving the quality of life of families by spacing and delaying the pregnancies and avoiding unwanted pregnancies. So it is recognized as a key life-saving intervention for mothers and their children. <sup>(1)</sup>

FP can reduce more than 30% of maternal deaths and 10% of child mortality if couples space their pregnancies more than 2 years apart. <sup>(2)</sup> If pregnancies are closely spaced (within the first year postpartum) it risks both mother and baby. It can cause preterm delivery, low birth weight and small for gestational age etc. <sup>(3)</sup> Also risk of child mortality is higher for very short birth-to-pregnancy interval (<12 months). If pregnancy interval was 24 months, under-five mortality would decrease by 13%. On the other hand, if couples wait 36 months, the decrease would be 25%. <sup>(4)</sup>

Postpartum family planning (PPFP) is the concept which mainly focuses on prevention of unintended and closely spaced pregnancies through the first 12 months following childbirth. <sup>(5)</sup> There are various methods which can be used as PPFP such as post-partum sterilization, Post-Partum Intra Uterine Device

(PPIUD), progesterone only methods (sub dermal implants and injections), Oral Contraceptive Pills and condoms. <sup>(5)</sup> According to an analysis of Demographic and Health Surveys data from 27 countries, 95% of women who are 0–12 month postpartum want to avoid a pregnancy in the next 24 months; but 70% of them are not using contraception. <sup>(6)</sup>

Health care personals can educate family regarding integrating PPFP opportunities during Ante Natal Clinic (ANC) visits, ante natal wards, post-natal wards and neonatal and child health clinics. <sup>(5)</sup> Also mothers can receive PPFP facilities within first 48 hours from where they deliver the baby.

Postpartum Intrauterine Device (PPIUD), Sub dermal Implants and Female Sterilization can be offered within first 48 hours after the delivery in tertiary care hospital. Ministry of Health Sri Lanka in collaboration with Sri Lanka College of obstetricians and gynecologists (SLCOG) implemented a project promoting PPIUD use within first 48 hours after delivery. <sup>(7)</sup>

Even though postpartum family planning facilities are available in majority of hospitals in Sri Lanka in both public and private sectors, there is a significant unmet need.

\*Corresponding author: A.M.R.K. Atapattu

Department of Obstetrics & Gynaecology, De Soysa Hospital For Women

Studies done in Sri Lanka showed the effectiveness on education about FP during antenatal clinic, antenatal wards and pre discharge is high.

## METHODOLOGY

### Study design

The present study was a descriptive cross-sectional study.

### Study Setting

The study was conducted in all four Obstetric units in De Soysa Hospital for Women, Colombo 08 (DSHW).

### Study Population

Inclusion criteria -Postpartum Mothers (within first 48 hours after delivery) who are delivering their babies in DMH in the time from 25<sup>th</sup> of September 2019 until the sample size is reached.

### Exclusion criteria

1. Mothers whose babies are in PBU or NICU.
2. Mothers who are having stillbirth in this pregnancy.
3. Mothers who are diagnosed to have postpartum psychological issues

### Sampling and sample size

The study will include all mothers in study population and the data was collected from 25th of September 2019 until the sample size is reached.

### Sampling: Expected proportion

From a primary survey done in De Soysa Hospital for Women in 2018, the acceptance of PFP in DSHW in 2017 was 25%. According to that the expected proportion for this research was taken as 0.25

$$\text{Sample size} = Z_{1-\alpha/2}^2 p(1-p)/d^2$$

$$= 1.96^2 \cdot 0.25 \cdot (1-0.25) / 0.05^2 = 288.12$$

$$\approx 288.12$$

Sample size = 289

With 10% of non-respondent rate – sample size is 318

### Data collection

Investigators were the Intern House Officers working in relevant postnatal wards in DSHW in 2018. They will go through the admission book of particular ward and will select the study population. After finding the mother, the investigator will explain the study objectives; importance and outcome of the study and written informed consent was obtained. Only the mothers who are willing to participate to the study was selected. Then the interviewer will question the participant and mark the relevant responses given by the mother.

### Data analysis

Date from all questionnaire was first entered into Microsoft excel sheets. Then they were analyzed using SPSS statistical analytical software.

### Ethical concerns

There are no major ethical issues related to this study; No interventions, invasive procedures were carried out. The target population is not a vulnerable group. The names and addresses will not be collected as a part of data collection. The data were stored in password-protected computers and data will only be

accessible to the investigators. Ethical clearance was obtained from Ethics Review Committee, National Hospital of Sri Lanka.

### Administrative clearance

The administrative clearance was obtained from Director of DSHW.

## RESULTS

A descriptive cross-sectional study was conducted at all four Obstetric units in De Soysa Hospital for Women, Colombo 08 (DSHW) among 295 mothers.

Table 1 Demographic Information

Variable	Frequency (%)
<b>Age</b>	28 ± 5.59 years
<b>Religion</b>	
Buddhist	100 (33.9)
Islam	88 (29.8)
Hindu	69 (23.4)
Catholic	38 (12.9)
<b>Highest Educational Qualification of the mother</b>	
Less than O/L	101 (34.2)
O/L	87 (29.5)
A/L	85 (28.8)
Degree	22 (7.5)
<b>Highest Educational Qualification of the farther</b>	
Less than O/L	104 (35.4)
O/L	107 (36.4)
A/L	63 (21.4)
Degree	20 (6.8)
<b>Number of children</b>	
1	158 (53.7)
2	80 (27.2)
3	45 (15.3)
4	9 (3.1)
5	2 (0.7)
<b>Age of the last child</b>	5 ± 2.43 years
<b>Monthly income</b>	
Less than 25000	54 (18.6)
25000-50000	161 (55.5)
More than 50000	75 (25.9)




Figure 1 Age ditribution

Study was conducted among 295 mothers with average mean age of 28 ± 5.59 years with majority Buddhist 100 (33.9%) followed by Islam of 88 (29.8%) out of total sample. Considering the level of education of mothers, 101 (34.2%) had less than O/L qualification, 87 (29.5%) completed O/L and 85

(28.8) completed A/L, similarly for the fathers, 101 (34.2%) had less than O/L qualification followed by 107 (36.4%) whom completed O/L and 63 (21.4%) completed A/L.

With regards to the number of children, more than half of the sample 158 (53.7%) having least one child and 80 (27.2%) mentioned the fact that they have least two children in the family. The mean age of last child noted as 5 ± 2.43 years. Lastly on the income level of the participants, 161 (55.5%) of majority earned an income between 25,000 to 50,000LKR and 75 (25.9%) from total studied sample earning a monthly income more than 50,000LKR.


**Table 2** Associated factors

Variable	Frequency (%)
<b>Mode of delivery of current pregnancy</b>	
NVD	171 (58.0)
Instrumental delivery	10 (3.4)
EM/LSCS	74 (25.1)
EL/LSCS	40 (13.6)
<b>Acceptance of contraceptives</b>	
Yes	152 (51.5)
No	143 (48.5)
<b>Type of contraceptive accepted</b>	
PPIUD	48 (33.8)
Sub dermal implant	82 (57.7)
PP- LRT	12 (8.5)
<b>Whose advices lead for the acceptance</b>	
MOH	19 (12.7)
PHM	28 (18.0)
Antenatal classes in Hospital	48 (31.3)
Doctors in ANC	31 (20.7)
Doctors in Antenatal wards	19 (12.7)
Other	7 (4.7)
<b>Time of deciding the acceptance</b>	
Before pregnancy	6 (4.0)
During PMH home visit	24 (16.0)
During antenatal clinics	86 (57.3)
During antenatal ward stay	14 (9.3)
During first 48 hours of postpartum period	20 (13.3)
<b>Reason for acceptance</b>	
To prevent unwanted pregnancies	16 (10.95)
To space the pregnancies	81 (44.5)
To limit the family size	49 (33.6)
Other reasons	16 (10.95)
<b>Reason for not accepting</b>	
Religious belief	8 (5.7)
Wants to have many children	18 (12.9)
Husband disapproval	31 (20.1)
Fear of side effects	20 (12.1)
Husband is not in Sri Lanka	11 (7.8)
Hope to use in future- Interval methods	53 (37.8)
Other reasons	5 (3.6)

Vaginal deliveries among 171 (58.0%) mothers resulted as the prominent mode of delivery of current pregnancy followed by EM/LSCS which managed among 74 (25.1%) mothers and more than half of the sample, 152 (51.5%) accepted the contraceptives. Considering the types of contraceptives accepted, a significantly higher proportion accepted Subdermal implant and 48 (33.8%) mothers accepted PPIUD type as contraceptive type.

Antenatal classes in Hospital of 48 (31.3%) and doctors in ANC of 31 (20.7%) detected as widely preferable advising sourced which lead for the acceptance and 86 (57.3%) of mothers deciding the acceptance during antenatal clinics followed by 24 (16.0%) mothers who accepting during PMH home visit. To keep space between pregnancies mentioned as the most common fact to accept the contraceptives and to limit the family size secondly commonest reason for acceptance. Nevertheless, 53 (37.8%) mothers mentioned hope to use in

future- Interval methods as the fact to not to accept particular management method along with 31 (20.1%) who do not have husband approval to proceed with the contraceptive method.



**Figure 2** Types of interval contraceptive methods

Among the mothers accepted the interval contraceptive methods, Sub dermal Implants considered as the most preferred with 29 (42.6%) mothers and DMPA Injections noted as next highly accepted method among 27 (39.7%) mothers.

**Table 3** Demographic factors associated with contraceptive methods

Variable	Hope to use in future (N=53)	Specific Reason (N=93)	P Value
<b>Age</b>			
Less than 30 Years	43 (81.1)	64 (68.8)	0.106
More than 30 Years	10 (18.9)	29 (31.2)	
<b>Religion</b>			
Buddhist	23 (43.4)	29 (31.2)	0.015
Islam	15 (28.3)	35 (37.6)	
Hindu	6 (11.3)	24 (25.8)	
Catholic	9 (17.0)	5 (5.4)	
<b>Highest Educational Qualification of the mother</b>			
Up to O/L	34 (64.2)	61 (65.6)	0.861
Above O/L	19 (35.8)	32 (44.4)	
<b>Highest Educational Qualification of the farther</b>			
Up to O/L	37 (69.8)	71 (77.2)	0.327
Above O/L	16 (30.2)	21 (22.8)	
<b>Number of children</b>			
One	30 (56.6)	62 (66.7)	0.105
Two	19 (35.8)	19 (20.4)	
Three Or More	4 (7.5)	12 (12.9)	
<b>Age of the last child</b>			
Up to 4 Year	7 (30.4)	12 (38.7)	0.529
Above 4 Year	16 (69.6)	19 (61.3)	
<b>Monthly income</b>			
Less than 25000	9 (17.0)	11 (12.1)	0.598
25000-50000	32 (60.4)	54 (59.3)	
More than 50000	12 (22.6)	26 (28.6)	

About the acceptance of the contraceptives, there is a significant association observed among religion of the participants and preferability (p=0.015)

**Table 4** Hypothesis testing

Hypothesis	P Value	Decision	Conclusion
H1: There is relationship between Reason for not accepting and Age	0.106	Do not reject HO	There is not a significance relationship between Reason for not accepting and Age
H2: There is relationship between Reason for not accepting and Religion	0.015	HO reject	There is a significance relationship between Reason for not accepting and Religion
H3: There is relationship between Reason for not	0.861	Do not reject HO	There is not a significance

accepting and Educational Qualification of the Mother			relationship between Reason for not accepting and Educational Qualification of the Mother
H4: There is relationship between Reason for not accepting and Educational Qualification of the farther	0.327	Do not reject HO	There is not a significance relationship between Reason for not accepting and Educational Qualification of the farther
H5: There is relationship between Reason for not accepting and Number of children	0.105	Do not reject HO	There is not a significance relationship between Reason for not accepting and Number of children
H6: There is relationship between Reason for not accepting and Age of the last child	0.529	Do not reject HO	There is not a significance relationship between Reason for not accepting and Age of the last child
H7: There is relationship between Reason for not accepting and Monthly income	0.598	Do not reject HO	There is not a significance relationship between Reason for not accepting and Monthly income

## DISCUSSION

Postpartum family planning (PPFP) is an essential service to prevent unwanted pregnancies, as well as to keep birth spacing for at least two years of duration, thus preventing poor maternal, perinatal, and neonatal health outcomes in subsequent pregnancies (1,2). Previous surveys reported regarding the unmet need for Postpartum family planning (PPFP) especially from the developing world, because many women become sexually active within the postpartum period prior initiating a contraceptive method<sup>(3)</sup>. Therefore, PPFP has been a critical component of reproductive health as stated by the WHO (4). Women's contraceptive preference will change following childbirth during the postpartum period of six weeks if the woman is lactating and in regards with non-lactating mother's postpartum period can be altered as 3 or 4 weeks<sup>(2)</sup>.

Our study revealed that about 152 (51.5 %) of the participants accepted postpartum family planning (PPFP) while 143 (48.5%) study participants denied PPFP. Moreover, only 20 (13.3 %) of participants accepted contraceptive methods during the first 48 hours of the postpartum period. A study conducted in a tertiary hospital, Thamilnadu, to rule out the readiness and admissibility of postpartum intrauterine contraceptive devices showed that 82.8 % primiparous mothers accepted PPFP (5). This percentage is quite higher than our results. In contrast, a study conducted in south-east Ethiopia by Alemayehu et al reported a low percentage, around 12.4% women accepted PPFP and this low percentage was due to religious beliefs and husband refusal<sup>(6)</sup>. These differences might be explained by the cultural impact and different degree of involvement of government and non-governmental reproductive health care organizations.

Considering the reasons for not accepting the PPFP among the women participated in our study, most of our participants 53 (37.8%) were hoping to use in future (Interval methods) followed by 31 (20.1 %) husband's disapproval, 20 (12.1%) fear of side effects and 18 (12.9%) wanted to have many

children. Similarly, Ethiopian study showed that commonest causes for rejection of PPFP was fear of complications (24.8%), religious beliefs (19.8%), and husband refusal (17.7%)<sup>(6)</sup>. Factors leading to not accepting PPFP might be associated with the educational level of husband and wife as well as awareness regarding the contraceptive methods<sup>(7)</sup>. In contrast, our study does not show any relationship between educational qualification of the mother or husband as a reason for not accepting PPFP. Furthermore, we did not find factors such as number of children, age of the last child and monthly income as reasons for not accepting PPFP. However, there is a significant association observed among the religion of the participants.

The most popular PPFP method among our participants was Sub dermal implant 82 (57.7 %), followed by PPIUD in 48 (33.8%) participants and leastly PP- LRT among 12 (8.5%) participants. The reason for Sub dermal implant to become the most preferable PPFP might be due to being a long term contraceptives user need minimum attention, low percentage of contraceptive failure and fertility is reversible after removal of implant according to their wish<sup>(8)</sup>. It is found that sub dermal implants are the commonest [n=29 (42.6%)] interval contraceptive method among our participants followed by DMPA Injections among 27 (39.75%) participants. A meta-analysis done by *Rubee et al* demonstrated that Long-acting reversible contraceptive usage has increased dramatically in African region followed by barrier methods, while IUCD use has also risen slightly<sup>(9)</sup>.

## CONCLUSION & RECOMMENDATION.

In this study, slightly above 50% of the study population only accepted the postpartum family planning method and subdermal implant is the commonest mode of contraception. The commonest factor for denying contraception was expectation of using Interval contraceptive method in future. However, more awareness programs should be carried out to encourage postpartum women in adhering to PPFP to prevent unwanted pregnancies, hence improving the maternal and infant quality of life.

## References

1. Who WHO. World Health Statistics [Internet]. Vol. 27, WHO World Health Organization. 2012.171p. Available from: [http://books.google.com/books?hl=en&amp;lr=&id=osms0oMN\\_qwC&oi=fnd&pg=PA7&dq=World+Health+Statistics&ots=iZU1J3AqjI&sig=sIBAWCK7qn9DYyQU\\_cHVERTK4Ks](http://books.google.com/books?hl=en&amp;lr=&id=osms0oMN_qwC&oi=fnd&pg=PA7&dq=World+Health+Statistics&ots=iZU1J3AqjI&sig=sIBAWCK7qn9DYyQU_cHVERTK4Ks)
2. Cleland J, Bernstein S, Ezeh A, Faundes A, Glasier A, Innis J. Family planning: the unfinished agenda. *Lancet* [Internet]. 2006 Nov 18 [cited 2018 Jan 1];368(9549):1810–27. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17113431>
3. DaVanzo J, Hale L, Razzaque A, Rahman M. Effects of interpregnancy interval and outcome of the preceding pregnancy on pregnancy outcomes in Matlab, Bangladesh. *BJOG* [Internet]. 2007 Sep [cited 2018 Jan 1];114(9):1079–87. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17617195>
4. Fotso JC, Cleland J, Mberu B, Mutua M, Elungata P. Birth spacing and child mortality: an analysis of prospective data from the Nairobi urban health and demographic surveillance system. *J Biosoc Sci* [Internet].

- 2013 Nov [cited 2018 Jan 1];45(6):779–98. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22958417>
5. WHO. Programming Strategies for Postpartum Family Planning. WHO Libr Cat Data Program. 2013;4.
  6. Pasha O, Goudar SS, Patel A, Garces A, Esamai F, Chomba E, et al. Postpartum contraceptive use and unmet need for family planning in five low-income countries. *Reprod Health* [Internet]. 2015 [cited 2018 Jan 1];12 Suppl 2(Suppl 2):S11. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26063346>
  7. SLCOG. PPIUD Project-Sri Lanka. [cited 2018 Mar 7]; Available from: [http://www.slkog.lk/img/PPIUD/News letterPPIUD Sri Lanka.pdf](http://www.slkog.lk/img/PPIUD/News%20letterPPIUD%20Sri%20Lanka.pdf)
  8. Who WHO. Health situation trends/ SEARO/Sri Lanka.
  9. Navodani KT, Fonseka P, Goonewardena CS. Postpartum Family Planning: Missed Opportunities across the Continuum of Care. *Ceylon Med J* [Internet]. 2017 Jun 30 [cited 2018 Mar 11];62(2):87. Available from:<http://www.ncbi.nlm.nih.gov/pubmed/28697589>
  10. Agampodi SB, Agampodi TC, Chandrasekara P. Family planning prevalence among postpartum mothers attending child welfare clinics - a sri lankan experience. *Indian J Community Med* [Internet]. 2009 Jul [cited 2018 Mar 11];34(3):265–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20049312>
  11. Yilmazel G, Balci E. Preferences and related factors for postpartum contraception in pregnant women. *Iran J Reprod Med* [Internet]. 2013 Oct [cited 2018 Mar 11];11(10):801–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24639700>
  12. Rao BS, Mathada VC. Awareness and Determinants of Contraceptive use among Nursing Mothers in Bellary, Karnataka. *J Clin DIAGNOSTIC Res* [Internet]. 2016 Jan [cited 2018 Mar 11];10(1):QC15-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26894128>
  13. Apanga PA, Adam MA. Factors influencing the uptake of family planning services in the Talensi District, Ghana. *Pan Afr Med J* [Internet]. 2015 [cited 2018 Mar 11];20. Available from: <http://www.panafrican-med-journal.com/content/article/20/10/full/>
  14. AT N, D G, G T. Postpartum Family Planning Utilization and Associated Factors among Women who Gave Birth in the Past 12 Months, Kebribeyah Town, Somali Region, Eastern Ethiopia. *J Women’s Heal Care* [Internet]. 2016 Dec 23 [cited 2018 Mar 11];5(6):1–12. Available from: <http://www.omicsgroup.org/journals/postpartum-family-planning-utilization-and-associated-factors-among-womenwho-gave-birth-in-the-past-12-months-kebribeyah-town-soma-2167-0420-1000340.php?aid=82753>
  15. Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Effects of birth spacing on maternal health: a systematic review. *Am J Obstet Gynecol*. 2007; 196(4): 297-308.
  16. Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Birth spacing and risk of adverse perinatal outcomes: a meta-analysis. *Jama*. 2006;295(15):1809–23.
  17. Borda MR, Winfrey W, McKaig C. Return to sexual activity and modern family planning use in the extended postpartum period: an analysis of findings from seventeen countries. *Afr J Reprod Health*. 2010;14(4 Spec):72–79. [http://www.who.int/reproductivehealth/publications/family\\_planning/ppfp\\_strategies/en/](http://www.who.int/reproductivehealth/publications/family_planning/ppfp_strategies/en/)
  18. Nadu T A cross sectional study on willingness and acceptability: PPIUCD by primipara mothers at a tertiary care hospital. 2017.
  19. Gonie A, Worku C, Assefa T, Bogale D, Girma A. Acceptability and factors associated with post-partum IUCD use among women who gave birth at bale zone health facilities Southeast-Ethiopia. *Contracept Reprod Med*. 2018;3:16. Published 2018 Nov 6. doi:10.1186/s40834-018-0071-z
  20. Hauck B, Costescu D. Barriers and Misperceptions limiting widespread use of intrauterine contraception. *J Obstet Gynaecol Canada*. 2015;37(7):606–616. doi: 10.1016/S1701-2163(15)30198-5.
  21. Peralta O, Diaz S, Croxatto H. Subdermal contraceptive implants. *J Steroid Biochem Mol Biol*. 1995 Jun;53(1-6):223-6.
  22. Dev, R., Kohler, P., Feder, M. et al. A systematic review and meta-analysis of postpartum contraceptive use among women in low- and middle-income countries. *Reprod Health* 16, 154 (2019). <https://doi.org/10.1186/s12978-019-0824-4>

\*\*\*\*\*