



ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

*International Journal of Recent Scientific Research*  
Vol. 6, Issue, 6, pp.4505-4507, June, 2015

**International Journal  
of Recent Scientific  
Research**

## RESEARCH ARTICLE

# A STUDY ON THE ESTIMATION OF HARDNESS IN GROUND WATER SAMPLES BY EDTA TRITRIMETRIC METHOD

**P.Ramya\*, A.Jagadeesh Babu<sup>1</sup>, E.Tirupathi Reddy<sup>2</sup> and L.Venkateswara Rao<sup>3</sup>**

<sup>1,3</sup>Department of Veterinary Public Health and Epidemiology College of Veterinary Science, Proddatur YSR Kadapa District, Andhrapradesh,  
Department of Poultry Science, College of Veterinary Science, Proddatur YSR Kadapa district, Andhrapradesh

### ARTICLE INFO

#### Article History:

Received 14<sup>th</sup>, May, 2015  
Received in revised form 23<sup>th</sup>,  
May, 2015  
Accepted 13<sup>th</sup>, June, 2015  
Published online 28<sup>th</sup>, June, 2015

#### Key words:

EDTA tritrimetric method, Groun water, Hardness.

### ABSTRACT

Water with high concentration of minerals is hard water. Water is essential for life. But water with very high degrees of hardness is harmful to health. 120 samples of ground water samples were collected from one town and two different villages. Hardness of water can be determined by EDTA tritrimetric method. Out of all the samples tested 39 (32.5%) samples were moderately hard, 76 (63.33%) samples were hard water and 5 (4.16%) samples were very hard water. Extreme degrees of hardness is dangerous to health. The present study did not revealed any soft water. There is a false notion that hard water is harmful to health, its not hard water, its very hard water (>180ppm). The minerals that are supplemented to body through water will be beneficial to health in several ways. So public should be educated about degrees of hardness and its effects.

**Copyright © P.Ramya et al** This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

### INTRODUCTION

Hardness, a physico chemical property of water, is generally a measure of calcium and magnesium ions in water. Zinc, iron, strontium, aluminum, and manganese can also contribute to water hardness; however, they are generally present in very low concentrations (NRC, 1974). These ions enter a water supply by leaching from minerals of rocks and soil. Common calcium-containing minerals are limestone (calcium carbonate) and chalk (calcium sulphate). A common magnesium mineral is dolomite, which also contains calcium (Gumashta et al., 2012). Initially, water hardness was understood to be a measure of the capacity of water to precipitate soap, which is in practice the sum of concentrations of all polyvalent cations present in water (Ca, Mg, Sr, Ba, Fe, Al, Mn, etc.) later it has been generally accepted that hardness is defined as the sum of the Ca and Mg concentrations, determined by the EDTA titrimetric method, and expressed in mmol/l (ISO, 1984) or as CaCO<sub>3</sub> equivalent in mg/l (Standard Methods, 1998), less frequently as the CaO equivalent (Kozisek, 2003). The type of anion found in these salts distinguishes between the two types of hardness - carbonate and non-carbonate hardness. Carbonate hardness is sometimes called temporary hardness because it can be removed by boiling water. Non-carbonate hardness cannot be broken down by boiling the water, so it is also known as

permanent hardness. In general, it is necessary to distinguish between the two types of hardness because the removal method differs for the two. Total hardness includes both temporary and permanent hardness caused by the calcium and magnesium, on the basis of which water is categorized as soft or hard and very hard (Sengupta, 2013). From the technical point of view, multiple different scales of water hardness were suggested (e.g. very soft – soft – medium hard – hard – very hard). Expectedly, both extreme degrees (i.e. very soft and very hard) are considered as undesirable concordantly from the technical and health points of view, but the optimum Ca and Mg water levels are not easy to determine since the health requirements may not coincide with the technical ones (Kozisek, 2003).

Health significance of water hardness was directly evidenced in the late 1950's. The relationship between water hardness and the incidence of vascular diseases was first described by a Japanese chemist Kobayashi (Kobayashi, 1957) who showed, based on epidemiological analysis, higher mortality rates from cerebrovascular diseases (stroke) in the areas of Japanese rivers with more acid (i.e. softer) water compared to those with more alkaline (i.e. harder) water used for drinking purposes. The World Health Organization says that "there does not appear to be any convincing evidence that water hardness causes adverse health effects in humans" (WHO, 2003). In

\*Corresponding author: **P.Ramya**

Department of Veterinary Public Health and Epidemiology College of Veterinary Science, Proddatur YSR Kadapa District, Andhrapradesh,

fact, the United States National Research Council has found that hard water can actually serve as a dietary supplement for calcium and magnesium (NRC, 1974). Hard drinking water is generally not harmful to one's health,<sup>[1]</sup> but can pose serious problems in industrial settings, where water hardness is monitored to avoid costly breakdowns in boilers, cooling towers, and other equipment that handles water. In domestic settings, hard water is often indicated by a lack of suds formation when soap is agitated in water, and by the formation of lime scale in kettles and water heaters. Wherever water hardness is a concern, water softening is commonly used to reduce hard water's adverse effects. The livestock owners in the Mega animal health camp area, conducted by College of Veterinary Science, Proddatur, expressed doubt about the effects of hard water on animal health and production and degrees of hardness. Keeping in view of all these things the present study was carried out to estimate the amount of hardness in the ground water of those areas.

### MATERIALS AND METHODS

One Hundred and twenty samples (120) of ground water were collected from one town and two different villages while attending Mega Animal Health Camp organized by College of Veterinary Science, Proddatur, YSR KAdapa district, Andhrapradesh, India. All the water samples were collected aseptically in to sterilized screw capped glass bottles and brought to the laboratory. Hardness of all the water samples was tested by using EDTA-tritrimetric method by taking 50ml of water sample into a conical flask along with 100ml of ammonia buffer solution and 100-200-mg of Erichrome black-T indicator followed by titration with EDTA solution present in a burette. End point is noted down by changing of the water solution color from wine red to blue and expressed as CaCO<sub>3</sub> equivalent in mg/l (Standard Methods, 1998). Amount of Hardness in water is calculated by using the formula.

$$\text{Hardness as mg/l CaCO}_3 = \frac{\text{ml of EDTA solution used} \times 1000}{\text{Volume of water samples taken}}$$

### RESULTS AND DISCUSSION

The present study had revealed that out of three places, the two villages had shown high level of hardness compared to the town. In Proddatur town, out of 40 water samples collected, almost all the samples were moderately hard (42.5%) and hard water samples (52.5%). Only few samples were very hard water (5%) (Table). The results of Villages were different from the results of town. Forty each number of samples were collected from two villages.

In Sannapalli village, more number of samples had shown hardness between 150-300mg/l of CaCO<sub>3</sub> (67.5%) i.e hardwater. Some of the samples had shown moderate hardness (30%) and very few of them have shown extreme hardness (2.5%)(Table). Similar type of results were found in the study of samples of Chinnakuravaluru village also. The results were hard water-70%, moderately hard water - 25% and very hard water - 5%. Finally no soft water sample was found in all the 120 samples (Table).

Altogether out of 120 samples, most of the samples are hard water (63.33%). There are only 5samples (4.16%) which are found to be very hard water. According to [Kozisek, \(2003\)](#) both the extreme degrees of hardness are dangerous to human health i.e very hard and very soft water. The present study did not revealed presence of any soft water, but it had reported very hard water in about 4.16% of the samples which is negligible. Water intake of cattle and milk production were unaffected by water containing up to 290 ppm of hardness (NRC, 1974). So the water in the above three places may not decrease the water intake and milk production of cattle in the above said areas.

Hard drinking water is generally not harmful to one's health, ([WHO, 2003](#)) but can pose serious problems in industrial settings. Most of the people especially housewives dislike hard water because it does not lather well or does not taste good, but they may not be knowing that it may prolong their lives, and more especially their husbands ([Time, Medicine, 2011](#)).

**United States Geological Survey had classified the water into hard and soft water, as per the table below**

Classification	hardness in mg/L	hardness in mmol/L	hardness in dGH/°dH	hardness in gpg	hardness in ppm
Soft	0-60	0-0.60	0-3.37	0-3.50	less than 60
Moderately hard	61-120	0.61-1.20	3.38-6.74	3.56-7.01	60-120
Hard	121-180	1.21-1.80	6.75-10.11	7.06-10.51	120-180
Very hard	181	1.81	10.12	10.57	> 180

**Table showing the results for hardness of different ground water samples**

S.No.	Place of sample collection	No.of samples	Classification based on degree of hardness in mg/L	
			No. of samples showing different degree of hardness	Percentage of samples showing different degree of hardness
1.	Proddatur town	40	S - 0	S - 0
			MH- 17	MH-42.5
			H - 21	H - 52.5
			VH - 02	VH - 5
2.	Sannapalli village	40	S - 0	S - 0
			MH- 12	MH- 30
			H - 27	H - 67.5
			VH - 01	VH - 2.5
3.	Chinnakuravaluru village	40	S - 0	S - 0
			MH- 10	MH- 25
			H - 28	H - 70
			VH - 02	VH - 05

Softwater-S, Moderately Hard-MH, Hard water-H, Very Hard water-VH

Calcium, one of the components of hard water, can be protective because it makes water less corrosive and less likely to leach toxic trace minerals, such as cadmium and lead, out of metal pipes (Seelig, 1989). According to the U.S. National Academy of Sciences by 1977 there had been more than 50 studies, in nine countries, that had indicated an inverse relationship between water hardness and mortality from cardiovascular disease (Harold and Foster, 1994). Most of the scientists have indicated a negative statistical association of various types of cancer morbidity/mortality with the hardness of water and calcium (Yang, 1998). Some studies correlate domestic hard water usage with increased eczema in children (McNally, 1998, Miyake *et al.*, 2004 and Arnedo-Pena, 2007).

## CONCLUSION

The present study had revealed extreme degrees of hardness in only 4.16% of the samples which may not be harmful to the people. Although most of the people dislikes to use hard water, magnesium and calcium are having some protective effect on cardiovascular mortality, the evidence being debated and does not prove causality and also the drinking water is the source of calcium and magnesium intake which are essential for the body. Most of the people are in false opinion that hard water is harmful to health, it's not hard water, it's very hard water (extreme degrees of hardness). So it would be better if we bring awareness among the public.

## Reference

1. Arnedo-Pena, A., Bellido-Blasco, J., Puig-Barbera, J., Artero-Civera, A., Campos-Cruaños, J.B., Pac-Sa, M.R., Villamarín-Vázquez, J.L. and Felis-Dauder, C. 2007. Domestic water hardness and prevalence of atopic eczema in Castellon (Spain) school children. *Salud Pública De México.*, 492 (4): 295–301.
2. Drinking Water Hardwater Hardness Calcium Magnesium Scale Stained Laundry. Water-research.net. Retrieved 2013-01-26.
3. Gumashta, J., Gumashtab, R., Sadawarte, S.K. 2012. Hard water and heart: the story revisited. *IOSR Journal of Pharmacy and Biological Sciences.*, 1(1): 07-20.
4. Harold D.F. 1994. Groundwater and Human Health,"Groundwater Resources of British Columbia,

Ministry of Environment, Lands, and Parks and Environment Canada. 6.1-6.3.

5. ISO. 1984. International Standard ISO 6059. Water quality – Determination of the sum of calcium and magnesium – EDTA titrimetric method. International Organization for Standardization. Geneva.
6. Kobayashi, J. 1957. On geographical relations between the chemical nature of river water and death rate from apoplexy. *Ber. Ohara Inst landwirt. Biol.*, 11: 12-21.
7. Kožíšek, F. 2003. Health significance of drinking water calcium and magnesium. National Institute of Public Health. A review.
8. McNally, N.J., Williams, H.C., Phillips, D.R., Smallman-Raynor, M., Lewis, S., Venn, A. and Britton, J. 1998. Atopic eczema and domestic water hardness. *The Lancet.*, 352(9127): 527–531.
9. Miyake, Y., Yokoyama, T., Yura, A., Iki, M. and Shimizu, T. 2004. Ecological association of water hardness with prevalence of childhood atopic dermatitis in a Japanese urban area. *Environ Res*, **94** (1): 33–7.
10. National Research Council. 1974. Nutrients and toxic substances in water for livestock and poultry. National Academy Science, Washington, D.C.
11. Seelig M.S. 1989. Cardiovascular Consequences Of Magnesium Deficiency And Loss: Pathogenesis, Prevalence And Manifestations- Magnesium and Chloride Loss In Refractory Potassium Repletion. *The American Journal of Cardiology.*, 63(14): G4-G21.
12. Sengupta, P. 2013. Potential Health Impacts of Hard Water. *Int. J of Prev. Med.*, 4(8): 866-875.
13. Standard Methods for the Examination of Water and Wastewater (1998). – #2340 Hardness. 20th ed. American Public Health Association / American Water Works Association / Water Environment Federation, Washington, DC.
14. Time, Medicine: Hard Water, Soft Arteries? (Monday, May. 02, 1960) VOL LXXV NO.18 Available at <http://www.time.com/time/magazine/0,9263,7601600502,00.html> . accessed on December 30th, 2011.
15. World Health Organization. (2003). Hardness in drinking water.
16. Yang, C.Y. 1998. Calcium and magnesium in drinking water and risk of death from cerebrovascular disease. *Stroke.*, 29: 411-4.

### How to cite this article:

P.Ramya *et al.*, A Study On The Estimation Of Hardness In Ground Water Samples By Edta Titrmetric Method. *International Journal of Recent Scientific Research Vol. 6, Issue, 6, pp.4505-4507, June, 2015*

\*\*\*\*\*