



*International Journal Of*  
**Recent Scientific  
Research**


ISSN: 0976-3031  
Volume: 7(6) June -2016

**RISK FACTORS OF ACUTE CELLULITIS IN ADULT PATIENTS: A CASE-CONTROL  
STUDY**

Nassaji M., Ghorbani R and Ghashghaee S



THE OFFICIAL PUBLICATION OF  
INTERNATIONAL JOURNAL OF RECENT SCIENTIFIC RESEARCH (IJRSR)  
<http://www.recentscientific.com/> [recentscientific@gmail.com](mailto:recentscientific@gmail.com)



ISSN: 0976-3031

Available Online at <http://www.recentscientific.com>

International Journal of Recent Scientific Research  
Vol. 7, Issue, 6, pp. 11905-11908, June, 2016

**International Journal of  
Recent Scientific  
Research**

## Research Article

### RISK FACTORS OF ACUTE CELLULITIS IN ADULT PATIENTS: A CASE-CONTROL STUDY

Nassaji M<sup>1\*</sup>., Ghorbani R<sup>2</sup> and Ghashghaee S<sup>3</sup>

<sup>1</sup>Infectious diseases, Social Determinants of Health Research Center, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran

<sup>2</sup>Biostatistics, Social Determinants of Health Research Center, Department of Community Medicine, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran

<sup>3</sup>Students' Research Committee, Semnan University of Medical Sciences, Semnan, Iran

#### ARTICLE INFO

##### Article History:

Received 11<sup>th</sup> March, 2016

Received in revised form 14<sup>th</sup> April, 2016

Accepted 18<sup>th</sup> May, 2016

Published online 28<sup>th</sup> June, 2016

##### Key Words:

Cellulitis; Risk factor; Adult;  
Case-control study

#### ABSTRACT

Cellulitis is a common medical condition with different severity. The identification of preventable risk factors could reduce morbidity and mortality from this disease. The aim of this study was to identify the risk factors associated with cellulitis in adult patients. We conducted a case-control study of 102 adult subjects with cellulitis and the same number of controls who were admitted to two university affiliated hospitals. Data were obtained with a questionnaire and from examination of patients. Two groups were matched based on age ( $P=0.717$ ) and gender ( $p=0.661$ ). The most common site of cellulitis was lower extremities (56.9%). The pack-year smoking and duration of smoking was significantly higher in patients with cellulitis ( $P=0<001$ ). Logistic regression analysis showed that recent trauma (OR=55.8, 95% CI: 11.2-278.4,  $p<0.001$ ), history of recent surgery (OR=6.3, 95% CI:1.7-23.7,  $p=0.006$ ), body mass index (OR=1.38, 95% CI:1.22-1.57,  $p<0.001$ ) and diabetes mellitus (OR=3.02, 95% CI:1.17-7.81,  $p=0.022$ ) was significantly associated with development of cellulitis. No association was observed between corticosteroid use and risk of cellulitis ( $p=0.158$ ). Our study showed that the most important risk factors for cellulitis in adult include trauma, recent surgery, high body mass index, diabetes mellitus and smoking.

Copyright © Nassaji M., Ghorbani R and Ghashghaee S., 2016, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Cellulitis is an inflammatory skin condition with an infectious origin involving the dermis and subcutaneous tissues. It is a common medical condition with different severity varies from mild to life threatening disease (Gunderson CG, 2011). Most commonly, cellulitis involves the lower extremities but may involve any parts of the body (Swartz MN, 2004). The diagnosis of cellulitis is generally based on the morphologic features of the lesion and the clinical setting characterized by erythema, swelling, warmth, and pain. Fever and constitutional symptoms are often mild or absent in uncomplicated cellulitis (Bailey E *et al*, 2011).

Several risk factors for the cellulitis have been described. The most common are trauma, edema, superficial or deep venous insufficiency, prior radiation therapy, previous surgery and skin disorders that cause a potential portal of entry such as toe web intertrigo (Stevens DL *et al*. 2009). Some other potential risk factors reported in studies including diabetes mellitus, prior malignancy, body mass index (BMI), smoking, alcoholism and ethnicity with controversial findings. Most of these previous

studies reported that general risk factors have little role in the development of cellulitis (Björnsdóttir S *et al* 2005, Dupuy A *et al* 1999, Halpern J *et al* 2008, Mokni M *et al* 2006).

The better identification of preventable risk factors for cellulitis has significant implications for reducing morbidity and mortality from this diseases and improving patient management. In addition differences findings in previous studies have suggested that there may be regional variations in risk factors and authors proposed more studies. So that we conducted a case control study to assess risk factors of cellulitis, particularly the roles of systemic risk factors.

## MATERIALS AND METHODS

This case-control study was conducted from May 2013 to June 2014 in two university-affiliated hospitals in Semnan, Iran. Adult patients ( $\geq 18$  years) with diagnosis of acute cellulitis who admitted in hospitals or were visited outpatients considered for participation in the study. Control group were selected from adult outpatient and without history of cellulitis. Informed consent was obtained from all subjects before enrollment.

\*Corresponding author: Nassaji M

Infectious diseases, Social Determinants of Health Research Center, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran

Patients with chronic ulcer, abscess, necrotizing fasciitis and patients with a main diagnosis or a reason for hospitalization that was not cellulitis were excluded.

Cellulitis was defined based on clinical findings by a board-certified specialist in infectious diseases. The clinical criteria included of pain, warmth, edema and erythema without sharp demarcation from uninvolved skin with or without fever (Bailey E *et al*, 2011).

A structured data collection sheet was used to collate information about demographics, as well as local and systemic risk factors. Baseline data collected included age, gender, body mass index, smoking status, diabetes mellitus and congestive heart failure. History of recent trauma (< 1 month), regional surgery in recent three month, corticosteroid consumption, and previous attack of cellulitis was also obtained. Location of cellulitis, extremity edema, dermatosis and toeweb intertrigo, were detected by clinical examination.

Body mass index (BMI), defined as the weight in kilograms divided by the height in meters squared ( $\text{kg}/\text{m}^2$ ). Diabetes mellitus was defined as self-reported history of diabetes mellitus and use of oral hypoglycaemic agents or insulin. Subjects were defined as cigarette smokers if they had smoked 3 or more cigarettes per day for at least 6 months. Patients were followed for response to treatment and divided as  $\leq 3$  or  $> 3$  days responder.

The study protocol was approved by Research Council and Ethical Committee of the Semnan University of Medical Science.

Statistical analysis was performed by Kolmogorov-Smirnov, Chi-Square, Student's t, Mann-Whitney tests and logistic regression analysis using SPSS16.0. P-value less than 0.05 considered statistically significant.

## RESULTS

Of all patients admitted with diagnosis of cellulitis that were screened 102 met inclusion criteria and were enrolled. Same numbers of controls were selected. The mean ( $\pm$  SD) age of patients was  $45.6 \pm 16.8$  years and the controls was  $46.5 \pm 16.8$  that showed no statistically significant difference ( $P=0.717$ ). 62.7% of cases and 65.7% of controls were male with no significant difference ( $p=0.661$ ). Two groups were matched based on age and gender. Baseline characteristics and co-morbidities of two groups are showed in table 1.

The infection was located in the lower extremities in most the patients (table 2). Nineteen (18.6%) of patients had fever on admission. Eleven of the cellulite cases (10.8%) had previous episodes of cellulitis. Duration of response to treatment was  $\leq 3$  days in 28.4% and  $> 3$  days in 71.6%.

From all patients 38.2% had history of trauma, whereas only 2% of control have positive history ( $p=0.001$ ). Mean BMI ( $\pm$ SD) of patients was  $25.8 \pm 4.1$  and for control was  $22.5 \pm 4.1$   $\text{Kg}/\text{m}^2$ . BMI of patients with cellulitis was significantly higher than controls ( $p<0.001$ ).

History of corticosteroid use was positive in 6.9% of patients and 12.7% of controls ( $p=0.158$ ). Mean ( $\pm$ SD) duration of corticosteroid use was  $8.7 \pm 8.0$  and  $2.4 \pm 8.0$  in patients and

controls respectively ( $p=0.056$ ). From patients 27.5% and from control 13.7% had diabetes mellitus that was significantly difference ( $p=0.015$ ).

Among the cases, 24 (24.5%) of subjects were current regular cigarette smokers whereas in control group 19 (18.6%) were smoker. The mean ( $\pm$  SD) number of pack-year smoking was  $16.4 \pm 20.3$  in cases and  $3.7 \pm 2.6$  in control group that was significantly different ( $P<0.001$ ). The mean ( $\pm$  SD) duration of smoking was  $15.21 \pm 9.9$  and  $7.0 \pm 3.8$  years in patients and control subjects respectively. There was significant difference in years of smoking between smoker subjects with and without cellulitis ( $P=0.001$ ).

Logistic regression analysis showed that BMI (OR=1.38, 95% CI: 1.22-1.57,  $p<0.001$ ), diabetes mellitus (OR=3.02, 95% CI: 1.17-7.81,  $p=0.022$ ), history of recent trauma (OR=55.8, 95% CI: 11.2-278.4,  $p<0.001$ ) and recent surgery (OR=6.3, 95% CI: 1.7-23.7,  $p=0.006$ ), was significantly associated with development of cellulitis (table 3).

**Table 1** Basic characteristics of case with cellulitis and control groups

Characteristic	Study Group			
	Cellulites (n=102)		Control (n=102)	
	n	%	n	%
Age(years)				
<30	20	19.6	20	19.6
30-39	19	18.6	19	18.6
40-49	24	23.5	23	22.5
50-59	18	17.6	17	16.7
60-69	12	11.8	14	13.7
$\geq 70$	9	8.8	9	8.8
BMI <sup>a</sup> (Kg/m <sup>2</sup> )				
<25	45	44.1	84	82.4
25-29.9	37	36.3	17	16.7
$\geq 30$	20	19.6	1	1.0
Co-morbidities				
-	64	62.7	63	61.8
Diabetes mellitus	28	27.5	14	13.7
CHF <sup>b</sup>	8	7.8	13	12.7
Corticosteroid use	7	6.9	13	12.7

<sup>a</sup> Body Mass Index <sup>b</sup> Congestive heart failure

**Table 2** Anatomical location of cellulitis in patients

Site of infection	n	%
Lower extremities	58	56.9
Upper extremities	32	31.4
Thorax/trunk	5	4.9
Head/Face	5	4.9
Genital	2	2

**Table 3** Logistic regression analysis for risk factors of cellulites in adult patients

Characteristic	Odds Ratio(OR)	95% Confidence Interval for OR	p-value
BMI <sup>a</sup> (Kg/m <sup>2</sup> )	1.38	1.22-1.57	<0.001
Co-morbidities			
-	1	-	-
Diabetes mellitus	3.02	1.17-7.81	0.022
CHF <sup>b</sup>	1.07	0.31-3.73	0.916
Pack- year smoking	1.13	1.00-1.29	0.056
Recent trauma	55.8	11.2-278.4	<0.001
Surgery	6.3	1.7-23.7	0.006

<sup>a</sup> Body Mass Index <sup>b</sup> Congestive heart failure

## DISCUSSION

The role of some predisposing factors of cellulitis has not yet been elucidated. The prevalence of predisposing factors may vary between patient populations, and this may explain discordant findings among studies.

The anatomical distribution of cellulitis in our patients was similar to the most previous studies that reported most common site of involvement was the lower extremity (Koutica P *et al* 1999, Lazzarini L *et al* 2005, Pavlotsky F *et al* 2004, Perelló-Alzamora MR *et al* 2012, Krasagakis K *et al* 2010). Local factors especially disruption of the cutaneous barrier seemed to be the most important risk factors for cellulitis across studies. Previous trauma and regional surgery were significantly associated with cellulitis in our study. In accordance with our findings some previous studies reported that traumatic wound and previous surgery were risk factors for cellulitis (Karpelin M *et al* 2010, Roujeau JC *et al* 2004, Inghammar M *et al* 2014). Defects in cutaneous barrier caused by trauma and surgery facilitate the entry of pathogens to skin and development of infection. However, in a study participants with cellulitis had a lower incidence of recent trauma compared with controls (23% vs. 44%) (Björnsdóttir S *et al* 2005).

Our findings showed that higher BMI was significantly associated with cellulitis which is in line with the results of some earlier studies. Dupuy *et al* in their study reported that being overweight was a risk factor associated with cellulitis of the leg (Dupuy A *et al* 1999). Similarly, obesity and higher BMI was independent risk factors for acute cellulitis in some other studies (Mokni M *et al* 2006, Karpelin M *et al* 2010, Lewis SD *et al* 2006). In a prospective case-control study of 100 subjects with cellulitis and 200 control subjects, matched for age and sex, obesity was significantly more frequent in the case patient group, but being overweight was not (Björnsdóttir S *et al* 2005).

Overweight and obesity has been suggested to be a risk factor for some infections. Several epidemiological investigations and emerging data indicate obesity may increase infection susceptibility in clinical settings (Falagas ME *et al*, 2006). It is mediated by the effect of obesity on the immune response through a variety of immune mediators, which leads to susceptibility to infections (Milner JJ *et al*, 2012).

Significant association between BMI and risk of cellulitis was not reported in some studies. To identify risk factors for acute lower limb cellulitis in the U.K. population, researcher found that systemic risk factors including obesity were not associated with cellulitis (Halpern J *et al* 2008). Teerachaisakul *et al* evaluated risk factors of cellulitis in patients with lymphedema and reported that body mass index was not risk factor of cellulitis (Teerachaisakul M *et al* 2013).

In most previously published studies, diabetes was not reported as risk factors of cellulitis (Halpern J *et al* 2008, Lewis SD *et al*, 2006, Teerachaisakul M *et al* 2013). In contrast, we found that diabetes is an independent systemic risk factor for cellulitis. Similarly, Mokni *et al* in their study reported that diabetes was associated with increased risk of cellulitis (Mokni M *et al* 2006). Two studies conducted among hospitalized adults to evaluate the factors that contribute to the development of cellulitis. Diabetes mellitus was established underlying

conditions in patients with cellulitis (Koutica P *et al* 1999, Serdar ZA *et al*, 2011). In a retrospective cohort study in Canada the risk ratios of having an infectious disease between those with and without diabetes were calculated. Rate of cellulitis was 1.81 time more common in patients with diabetes (Shah BR *et al*, 2003). Skin and soft tissue infections were significantly differences in the frequencies between diabetic and non-diabetic individuals in suaya *et al* study (Suaya JA *et al*, 2013).

Individuals with diabetes have a greater frequency and severity of infections. The reasons for this include abnormality in cell mediated immunity and phagocyte functions associated with hyperglycemia (Jackson LA, 2005, Gupta S *et al*, 2007). Also diabetes causes some abnormality in skin condition that leads to disruption of integrity of skin, abnormal proprioception and neuropathy that predisposes to cellulitis (Oumeish OY, 2008).

The difference in these findings might at least partly be explained by differences in the study design, patients' selection, number of samples and confounding variables.

The result of study shows that patients with cellulitis have had significantly more pack-years of smoking than controls. Also, duration of smoking was significantly higher in patients with cellulitis. In accordance with this, a case-control retrospective study was conducted to evaluate potential risk factors for development of recurrent cellulitis. Authors found that tobacco use was significant independent predictors of cellulitis (Lewis SD *et al*, 2006). In a study on patients who underwent abdominoplasty, postoperative wound infection was more common in smokers than non smoker. Both number and duration of smoking increased risk of infections (Araco A *et al*, 2008). In another study smoking was significant risk factors for surgical site infections in orthopedic patients (Jain RK *et al*, 2004).

The currently available literature suggests that smoking exerts deleterious effects on both morphological and functional aspects of the microcirculation of skin and compromise perfusion of skin. The mechanisms of this phenomenon include compromised endothelial-dependent vasorelaxation, thickening of the walls of arterioles, platelet aggregation and endothelial cell dysfunction (Lehr HA, 2007). In addition; collagen synthesis and the deposition of mature collagen in the extracellular matrix are reduced by smoking. This phenomenon leads to an imbalance between biosynthesis and degradation of dermal proteins (Thomsen SF *et al*, 2010). In contrast some other studies did not show any association between smoking and risk of cellulitis (Björnsdóttir S *et al* 2005, Halpern J *et al* 2008, Mokni M *et al* 2006).

Most studies about cellulitis did not evaluate the role of corticosteroid use. Corticosteroid therapy did not appear to increase the risk of cellulitis in our patients. Similarly Kumar *et al* reported that corticosteroid administration did not increase the risk of postsurgical wound infection in their patients (Kumar S *et al*, 2013).

Our findings are strengthened by the case-control study design, age and gender matched groups and inclusion of both inpatients and outpatients. There are a few limitations to our study that deserve mention. First, the eligible patient number is relatively

small in this study. Second, we assessed patients in university affiliated hospital, so that referral bias may be occurred.

## CONCLUSION

In this study we found that beside of local risk factors, general risk factors including diabetes mellitus, body mass index and smoking are possible risk factors for development of cellulitis. Therefore, it is recommended that protection against trauma, controlling of blood glucose levels, reducing body weight and smoking cessation may lead to reducing incidence of cellulitis. Further prospective studies with larger samples are needed to better identification the role of these risk factors in acute cellulitis.

## References

- Araco A, Gravante G, Sorge R, Araco F, Delogu D, Cervelli V(2008). Wound infections in aesthetic abdominoplasties: the role of smoking. *Plast Reconstr Surg.* 121(5):305e-310e.
- Bailey E, Kroshinsky D (2011). Cellulitis: diagnosis and management. *Dermatol Ther.* 24(2):229-39
- Björnsdóttir S, Gottfredsson M, Thórisdóttir AS, Gunnarsson GB, Ríkardsdóttir H, Kristjánsson M *et al* (2005). Risk factors for acute cellulitis of the lower limb: a prospective case-control study. *Clin Infect Dis.* 41(10):1416-22.
- Dupuy A, Benchikhi H, Roujeau JC, Bernard P, Vaillant L, Chosidow O *et al* (1993). Risk factors for erysipelas of the leg (cellulitis): case-control study. *BMJ.* 18(7198):1591-4.
- Falagas ME, Kompoti M (2006). Obesity and infection. *Lancet Infect Dis.* 6(7):438-46.
- Gunderson CG (2011). Cellulitis: definition, etiology, and clinical features. *Am J Med.* 124(12):1113-22.
- Gupta S, Koirala J, Khardori R, Khardori N (2007). Infections in diabetes mellitus and hyperglycemia. *Infect Dis Clin North Am.* 21(3):617-38, vii.
- Halpern J, Holder R, Langford NJ (2008). Ethnicity and other **risk factors** for acute lower limb **cellulitis**: a U.K.-based prospective case-control study. *Br J Dermatol.* 158(6):1288-92.
- Inghammar M, Rasmussen M, Linder A (2014). Recurrent erysipelas--risk factors and clinical presentation. *BMC Infect Dis.* 14:270.
- Jackson LA (2005). Evaluating diabetes mellitus as a risk factor for community-acquired infections. *Clin Infect Dis.* 41(3):289-90.
- Jain RK, Shukla R, Singh P, Kumar R (2014). Epidemiology and risk factors for surgical site infections in patients requiring orthopedic surgery. *Eur J Orthop Surg Traumatol.* [Epub ahead of print]
- Karppelein M, Siljander T, Vuopio-Varkila J, Kere J, Huhtala H, Vuento R *et al*(2010). Factors predisposing to acute and recurrent bacterial non-necrotizing cellulitis in hospitalized patients: a prospective case-control study. *Clin Microbiol Infect.* 16(6):729-34.
- Koutica P, Mylonakis E, Boyce J (1999). Cellulitis: evaluation of possible predisposing risk factors in hospitalized patients. *Diagn Microbiol Infect Dis.* 34:325-7.
- Krasagakis K, Valachis A, Maniatakis P, Krüger-Krasagakis S, Samonis G, Tosca AD(2010). Analysis of epidemiology, clinical features and management of erysipelas. *Int J Dermatol.* 49(9):1012-7.
- Kumar S, van Popta D, Rodrigues-Pinto R, Stephenson J, Mohammad S, Siddique I *et al*(2013). Risk factors for wound infection in surgery for spinal metastasis. *Eur Spine J.* [Epub ahead of print]
- Lazzarini L, Conti E, Tositti G, de Lalla F (2005). Erysipelas and cellulitis: clinical and microbiological spectrum in an Italian tertiary care hospital. *J Infect.* 51(5):383-9.
- Lehr HA (2000). Microcirculatory dysfunction induced by cigarette smoking. *Microcirculation.* 7: 367-84.
- Lewis SD, Peter GS, Gomez-Marín O, Bisno AL (2006). Risk factors for recurrent lower extremity cellulitis in a US Veterans medical center population. *Am J Med Sci.* 332: 304-7.
- Milner JJ, Beck MA (2012). The impact of obesity on the immune response to infection. *Proc Nutr Soc.* 71(2):298-306.
- Mokni M, Dupuy A, Denguezli M, Dhaoui R, Bouassida S, Amri M *et al*(2006). Risk factors for erysipelas of the leg in Tunisia: a multicenter case-control study. *Dermatology.* 212(2):108-12.
- Oumeish OY (2008). Skin disorders in patients with diabetes. *Clin Dermatol.* 26(3):235-42.
- Pavlotsky F, Amrani S, Trau H (2004). Recurrent erysipelas: risk factors. *J Dtsch Dermatol Ges.* 2(2):89-95.
- Perelló-Alzamora MR, Santos-Duran JC, Sánchez-Barba M, Cañueto J, Marcos M, Unamuno P (2012). Clinical and epidemiological characteristics of adult patients hospitalized for erysipelas and cellulitis. *Eur J Clin Microbiol Infect Dis.* 31(9):2147-52.
- Roujeau JC, Sigurgeirsson B, Korting HC, Kerl H, Paul C (2004). Chronic dermatomycoses of the foot as risk factors for acute bacterial cellulitis of the leg: a case-control study. *Dermatology.* 209(4):301-7.
- Serdar ZA, Akçay SŞ, Inan A, Dağlı O (2011). Evaluation of microbiological spectrum and **risk factors** of **cellulitis** in hospitalized patients. *Cutan Ocul Toxicol.* 30(3):221-4.
- Shah BR, Hux JE (2003). Quantifying the risk of infectious diseases for people with diabetes. *Diabetes Care.* 26(2):510-3.
- Stevens DL, Eron LL (2009). Cellulitis and soft-tissue infections. *Ann Intern Med.* 150(1):ITC11.
- Suaya JA, Eisenberg DF, Fang C, Miller LG (2013). Skin and soft tissue infections and associated complications among commercially insured patients aged 0-64 years with and without diabetes in the U.S. *PLoS One.* 8(4):e60057.
- Swartz MN (2004). Clinical practice. Cellulitis. *N Engl J Med.* 350(9):904-12.
- Teerachaisakul M, Ekataksin W, Durongwatana S, Taneepanichskul S (2013). Risk factors for cellulitis in patients with lymphedema: a case-controlled study. *Lymphology.* 46(3):150-6.
- Thomsen SF, Sørensen LT(2010). Smoking and skin disease. *Skin Therapy Lett.* 15: 4-7.

T.SSN 0976-3031



9 770976 303009 >